arXiv:1203.6494v2 [math.MG] 10 Apr 2012

HYPERBOLIC LAMBERT QUADRILATERALS AND QUASICONFORMAL MAPPINGS

MATTI VUORINEN AND GENDI WANG

Abstract. We prove sharp bounds for the product and the sum of two hyperbolic distances between the opposite sides of hyperbolic Lambert quadrilaterals in the unit disk. Furthermore, we study the images of Lambert quadrilaterals under quasiconformal mappings from the unit disk onto itself and obtain sharp results in this case, too.

Keywords. Hyperbolic quadrilateral, quasiconformal mapping
2010 Mathematics Subject Classification. 51M09(51M15)

1. Introduction

Given a pair of points in the closure of the unit disk $\mathbb{B}^2$, there exists a unique hyperbolic geodesic line containing them. Hyperbolic lines are simply sets of the form $C \cap \overline{\mathbb{B}^2}$ where C is a circle perpendicular to the unit circle, and the Euclidean diameters of $\mathbb{B}^2$. For a quadruple of four points $\{a, b, c, d\}$ in the closure of the unit disk we can draw these hyperbolic lines through each of the four pairs of points $\{a, b\}$, $\{b, c\}$, $\{c, d\}$, and $\{d, a\}$. If these hyperbolic lines bound a domain $D \subset \mathbb{B}^2$ such that the points $\{a, b, c, d\}$ are in the positive order on the boundary of the domain ∂D , then we say that the quadruple of points $\{a, b, c, d\}$ determines a hyperbolic quadrilateral $Q(a, b, c, d)$ and that the points a, b, c, d are its vertices. A hyperbolic quadrilateral with angles equal to $\pi/2, \pi/2, \pi/2, \phi\, (0 \le \phi < \pi/2)$, is called a hyperbolic *Lambert* quadrilateral [Be, p. 156], see Figure 1. Observe that one of the vertices of a Lambert quadrilateral may be on the unit circle, in which case the angle at that vertex is $\phi = 0$.

In this paper, we study bounds for the product and the sum of two hyperbolic distances between the opposite sides of hyperbolic Lambert quadrilaterals in the unit disk. Also, we consider the same product expression for the images of these hyperbolic Lambert quadrilaterals under quasiconformal mappings from the unit disk onto itself. In particular, we obtain similar results for ideal hyperbolic quadrilaterals, i.e. in the case when all the vertices are on the unit circle and all the angles are zero. This follows, because an ideal hyperbolic quadrilateral can be subdivided into four Lambert quadrilaterals.

For the formulation of our main results we introduce some notation – further notation will be given below in Section 2. Let $J^*[a, b]$ be the hyperbolic geodesic line with the end points a and b on the boundary of $\mathbb{B}^2$, and $J[a, b]$ be the hyperbolic geodesic segment with end points a and b when $a, b \in \mathbb{B}^2$.

Given two subsets A, B of $\mathbb{B}^2$ (or of the upper half-plane $\mathbb{H}^2$), let $d_\rho(A, B)$ denote the hyperbolic distance between them, defined as

$$d_\rho(A, B) = \inf_{\substack{x \in A \\ y \in B}} \rho(x, y) \,,$$

where $\rho(x, y)$ stands for the hyperbolic distance (2.4)(or (2.3),resp.).

We now formulate our main results.

File: lamb120410.tex, printed: 2013-3-1, 22.01

Theorem 1.1. *Let* $Q(v_a\,, v_b\,, v_c\,, v_d)$ *be a hyperbolic Lambert quadrilateral in* $\mathbb{B}^2$ *and let the quadruple of interior angles* $(\frac{\pi}{2}\,, \frac{\pi}{2}\,, \phi\,, \frac{\pi}{2})$, $\phi \in [0, \pi/2)$, *correspond to the quadruple* $(v_a\,, v_b\,, v_c\,, v_d)$ *of vertices. Let* $d_1 = d_\rho(J[v_a, v_d]\,, J[v_b, v_c])$, $d_2 = d_\rho(J[v_a, v_b]\,, J[v_c, v_d])$ *(see Figure 1), and let* $C = \mathrm{th}\rho(v_a, v_c) \in (0, 1]$. *Then*

$$d_1 d_2 \le (\mathrm{arth}\,(\frac{\sqrt{2}}{2}C))^2\,.$$

Here equality holds if and only if v_c *is on the bisector of the interior angle at* v_a.



FIGURE 1. A hyperbolic Lambert quadrilateral in $\mathbb{B}^2$

Theorem 1.2. *Let* $Q(v_a\,, v_b\,, v_c\,, v_d)$, d_1, d_2 *and* C *be as in Theorem 1.1. Let* $m = \sqrt{(2 - C^2)(3C^2 - 2)}$, $r_0 = \sqrt{\frac{1 - m/C^2}{2}}$ *and* $r_0' = \sqrt{1 - r_0^2}$.

(1) If $0 < C \le \sqrt{\frac{2}{3}}$, *then*

$$\mathrm{arth}C < d_1 + d_2 \le \mathrm{arth}(\frac{2\sqrt{2}C}{2 + C^2}).$$

Equality holds in the right-hand side if and only if v_c *is on the bisector of the interior angle at* v_a.

(2) If $\sqrt{\frac{2}{3}} < C < \sqrt{2(\sqrt{2} - 1)}$, *then*

$$\mathrm{arth}C < d_1 + d_2 \le \mathrm{arth}(\frac{C(r_0 + r_0')}{1 + C^2 r_0 r_0'}).$$

Equality holds in the right-hand side if and only if the interior angle between $J[v_a, v_b]$ *and* $J[v_a, v_c]$ *is* $\arccos r_0$ *or* $\arccos r_0'$.

(3) If $\sqrt{2(\sqrt{2} - 1)} \le C < 1$, *then*

$$\mathrm{arth}(\frac{2\sqrt{2}C}{2 + C^2}) \le d_1 + d_2 \le \mathrm{arth}(\frac{C(r_0 + r_0')}{1 + C^2 r_0 r_0'}).$$

Equality holds in the left-hand side if and only if v_c *is on the bisector of the interior angle at* v_a. *Equality holds in the right-hand side if and only if the interior angle between* $J[v_a, v_b]$ *and* $J[v_a, v_c]$ *is* $\arccos r_0$ *or* $\arccos r_0'$.

(4) If $C = 1$, *then*

$$d_1 + d_2 \geq \operatorname{arth}(\frac{2\sqrt{2}}{3}) .$$

Equality holds if and only if v_c *is on the bisector of the interior angle at* v_a.

In a Lambert quadrilateral, the angle ϕ is related to the lengths d_1, d_2 of the sides "opposite" to it as follows [Be, Theorem 7.17.1, p. 157]:

$$\operatorname{sh} d_1 \operatorname{sh} d_2 = \cos\phi.$$

See also the recent paper of A. Beardon and D. Minda [BM, Lemma 5]. The proof of Theorem 1.1 yields the following corollary, which provides a connection between d_1, d_2 and $C = \operatorname{th}\rho(v_a, v_c)$.

Corollary 1.3. *Let* C, d_1 *and* d_2 *be as in Theorem 1.1. Then*

$$\operatorname{th}^2 d_1 + \operatorname{th}^2 d_2 = C^2.$$

Corollary 1.4. *Let* $Q(a, b, c, d)$ *be an ideal hyperbolic quadrilateral in* B^2. *Let* $d_1 = d_\rho(J^*[a, d], J^*[b, c])$ *and* $d_2 = d_\rho(J^*[a, b], J^*[c, d])$ *(see Figure 2). Then*

$$d_1 d_2 \leq (2\log(\sqrt{2} + 1))^2$$

and

$$d_1 + d_2 \geq 4\log(\sqrt{2} + 1) .$$

In both cases equalities hold if and only if $|a, b, c, d| = 2$.



FIGURE 2. An ideal hyperbolic quadrilateral in $\mathbb{B}^2$

Remark 1.5. $|a, b, c, d| = 2$ *means that there exists a Möbius transformation* f *such that* $f(a) = 1$,$f(b) = i$,$f(c) = -1$,$f(d) = -i$, *see* (2.2).

Theorem 1.6. *Let $f : \mathbb{B}^2 \to \mathbb{B}^2$ be a K-quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$ and let $Q(v_a, v_b, v_c, v_d)$ be a quadrilateral in $\mathbb{B}^2$ and d_1, d_2, C be as in Theorem 1.1.*

Denote $D_1 = d_\rho(f(J[v_a, v_d]), f(J[v_b, v_c]))$ and $D_2 = d_\rho(f(J[v_a, v_b]), f(J[v_c, v_d]))$.

(1) If $0 < C \le \frac{e^2-1}{e^2+1} \approx 0.761594$, then

$$D_1 D_2 \le c(K)^2 (\operatorname{arth}(\frac{\sqrt{2}}{2} C))^{2/K}.$$

(2) If $\frac{e^2-1}{e^2+1} < C \le 1$, then let $r_C = \frac{1}{C}\frac{e^2-1}{e^2+1} \approx \frac{0.761594}{C}$ and

$$M_C = \frac{f_C(\sqrt{1-r_C^2})}{f_C(r_C)} > 1.$$

Let $r_C(K)$ be a constant which depends on K and C and satisfies the equation $Kf_C(r) = f_C(\sqrt{1-r^2})$ with $r_C(K) \in (r_C, 1)$, where $f_C(r)$ is the same as in Lemma 2.9(1) by taking $c = C$. Further, define

$$d(x, C) = \operatorname{arth}(Cx)(\operatorname{arth}(C\sqrt{1-x^2}))^{1/K}, 0 < x < 1.$$

Then

$$D_1 D_2 \le c(K)^2 \max\{d(r_C(K), C),\ (\operatorname{arth}(\frac{\sqrt{2}}{2} C))^{2/K}\}$$

if $K > M_C$ and

$$D_1 D_2 \le c(K)^2 \max\{d(r_C, C),\ (\operatorname{arth}(\frac{\sqrt{2}}{2} C))^{2/K}\}$$

if $1 \le K \le M_C$, where $c(K)$ is as in Lemma 4.2.

Corollary 1.7. *Let $f : \mathbb{B}^2 \to \mathbb{B}^2$ be a K-quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$ and let $Q(a, b, c, d)$ be a quadrilateral in $\mathbb{B}^2$ and d_1, d_2 be as in Corollary 1.4. Denote $D_1 = d_\rho(f(J^*[a, d]), f(J^*[b, c]))$ and $D_2 = d_\rho(f(J^*[a, b]), f(J^*[c, d]))$. Further denote $r_1 = \frac{2\sqrt{e}}{e+1} \approx 0.886819$ and*

$$M_1 = \frac{(e-1)(\log(\sqrt{e}+1) - \log(\sqrt{e}-1))}{\sqrt{e}} \approx 1.46618$$

and define $r_1(K)$ to be a constant which depends on K and satisfies the equation $Kf_1(r) = f_1(\sqrt{1-r^2})$ with $r_1(K) \in (r_1, 1)$, and where $f_1(r)$ is the same as in Lemma 2.9(1) by taking $c = 1$. With the notation

$$d(x) = \operatorname{arth}(x)(\operatorname{arth}(\sqrt{1-x^2}))^{1/K}, 0 < x < 1,$$

we have

$$D_1 D_2 \le c(K)^2 \max\{2^{1+1/K} d(r_1(K)),\ (2\log(\sqrt{2}+1))^2\}$$

if $K > M_1$ and

$$D_1 D_2 \le c(K)^2 \max\{2^{1+1/K} d(r_1),\ (2\log(\sqrt{2}+1))^2\}$$

if $1 \le K \le M_1$, where $c(K)$ is as in Lemma 4.2.

2. Preliminaries

It is assumed that the reader is familiar with basic definitions of geometric function theory and quasiconformal mapping theory, see e.g. [Be, V]. We recall here some basic information on hyperbolic geometry [Be].

The chordal distance is defined by

$$\begin{cases} q(x,y) = \frac{|x-y|}{\sqrt{1+|x|^2}\sqrt{1+|y|^2}}; & x \neq \infty \neq y, \\ q(x,\infty) = \frac{1}{\sqrt{1+|x|^2}}, & \end{cases} \tag{2.1}$$

for $x, y \in \overline{\mathbb{R}^2}$.

For an ordered quadruple a,b,c,d of distinct points in $\overline{\mathbb{R}^2}$ we define the absolute (cross) ratio by

$$|a,b,c,d| = \frac{q(a,c)q(b,d)}{q(a,b)q(c,d)}.$$

It follows from (2.1) that for distinct points $a,b,c,d \in \mathbb{R}^2$

$$|a,b,c,d| = \frac{|a-c||b-d|}{|a-b||c-d|}. \tag{2.2}$$

The most important property of the absolute ratio is Möbius invariance, see [Be, p.72, Theorem 3.2.7], i.e. if f is a Möbius transformation, then

$$|f(a),f(b),f(c),f(d)| = |a,b,c,d|,$$

for all distinct a,b,c,d in $\overline{\mathbb{R}^2}$.

For a domain $G \subsetneq \mathbb{R}^2$ and a continuous weight function $w : G \to (0,\infty)$, we define the weighted length of a rectifiable curve $\gamma \subset G$ to be

$$l_w(\gamma) = \int_\gamma w(z)|dz|$$

and the weighted distance between two points $x, y \in G$ by

$$d_w(x,y) = \inf_\gamma l_w(\gamma),$$

where the infimum is taken over all rectifiable curves in G joining x and y. It is easy to see that d_w defines a metric on G and (G, d_w) is a metric space. We say that a curve $\gamma : [0,1] \to G$ is a geodesic joining $\gamma(0)$ and $\gamma(1)$ if for all $t \in (0,1)$, we have

$$d_w(\gamma(0),\gamma(1)) = d_w(\gamma(0),\gamma(t)) + d_w(\gamma(t),\gamma(1)).$$

The hyperbolic distance in $\mathbb{H}^2$ and $\mathbb{B}^2$ is defined in terms of the weight functions $w_{\mathbb{H}^2}(x) = 1/x_2$ and $w_{\mathbb{B}^2}(x) = 2/(1-|x|^2)$, resp. We also have the corresponding explicit formulas

$$\cosh \rho_{\mathbb{H}^2}(x,y) = 1 + \frac{|x-y|^2}{2x_2y_2} \tag{2.3}$$

for all $x, y \in \mathbb{H}^2$ [Be, p.35] and

$$\mathrm{th}\frac{\rho_{\mathbb{B}^2}(x,y)}{2} = \frac{|x-y|}{\sqrt{|x-y|^2 + (1-|x|^2)(1-|y|^2)}} \tag{2.4}$$

for all $x, y \in \mathbb{B}^2$ [Be, p.40]. In particular, for $t \in (0,1)$,

$$\rho_{\mathbb{B}^2}(0, te_1) = \log\frac{1+t}{1-t} = 2\mathrm{arth}t. \tag{2.5}$$

There is a third equivalent way to express the hyperbolic distances. Let $G \in \{\mathbb{H}^2, \mathbb{B}^2\}$, $x, y \in G$ and let L be an arc of a circle perpendicular to ∂G with $x, y \in L$ and let $\{x_*, y_*\} = L \cap \partial G$, the points being labelled so that x_*, x, y, y_* occur in this order on L. Then by [Be, p.133,(7.26)]

$$\rho_G(x,y) = \sup\{\log|a,x,y,b| : a,b \in \partial G\} = \log|x_*,x,y,y_*|. \tag{2.6}$$

We will omit the subscript G if it is clear from the context. The hyperbolic distance is invariant under Möbius transformations of $\mathbb{B}^2$ onto $\mathbb{B}^2$ or onto $\mathbb{H}^2$.

Hyperbolic geodesics are arcs of circles which are orthogonal to the boundary of the domain. More precisely, for $a, b \in \mathbb{B}^2$ (or $\mathbb{H}^2$), the hyperbolic geodesic segment joining a to b is an arc of a circle orthogonal to S^1 (or $\partial\mathbb{H}^2$). In a limiting case the points a and b are located on a Euclidean line through 0 (or located on a normal of $\partial\mathbb{H}^2$), see [Be]. Therefore, the points x_* and y_* are the end points of the hyperbolic geodesic. For any two distinct points the hyperbolic geodesic segment is unique (see Figure 3 and 4). For basic facts about the hyperbolic geometry we refer the interested reader to [A], [Be] and [KL].



Figure 3. Hyperbolic geodesic segments in $\mathbb{H}^2$



Figure 4. Hyperbolic geodesic segments in $\mathbb{B}^2$

By [K, p.6, Exercise 1.1.27] and [KV, Lemma 2.2], for $x, y \in \mathbb{R}^2 \setminus \{0\}$ such that $0, x, y$ are noncollinear, the circle $S^1(a, r_a)$ containing x, y is orthogonal to the unit circle, where

$$a = i\frac{y(1+|x|^2) - x(1+|y|^2)}{2(x_2y_1 - x_1y_2)} \quad and \quad r_a = \frac{|x-y|\big|x|y|^2 - y\big|}{2|y||x_1y_2 - x_2y_1|}. \tag{2.7}$$

For $r, s \in (0, +\infty)$, the *Hölder mean of order p* is defined by

$$H_p(r,s) = \left(\frac{r^p + s^p}{2}\right)^{1/p} \quad \text{for} \quad p \neq 0, \quad H_0(r,s) = \sqrt{r\,s}.$$

For $p = 1$, we get the arithmetic mean $A = H_1$, for $p = 0$ the geometric mean $G = H_0$, and for $p = -1$ the harmonic mean $H = H_{-1}$. It is well-known that $H_p(r,s)$ is continuous and increasing with respect to p. Many interesting properties of Hölder means are given in [Bu] and [HLP].

A function $f : I \to J$ is called $H_{p,q}$*-convex(concave)* if it satisfies

$$f(H_p(r,s)) \leq (\geq) H_q(f(r), f(s))$$

for all $r, s \in I$, and *strictly* $H_{p,q}$*-convex(concave)* if the inequality is strict except for $r = s$. For $H_{p,q}$-convexity of some special functions the reader is referred to [AVV2, Ba, ZWC].

Some other notation are also needed in the paper. Let $[a, b]$ be the Euclidean segment with end points a and b. Let X, Y be the real axis and imaginary axis, respectively. Let $\mathrm{Arc}(abc)$ be the circular arc with end points a, c and through b and $\mathrm{SArc}(ac)$ be semicircle with end point a, c.

The so-called *monotone form of l'Hôpital's rule* is useful in deriving monotonicity properties and obtaining inequalities.

Lemma 2.8. [AVV1, Theorem 1.25]. *For* $-\infty < a < b < \infty$*, let* $f, g : [a, b] \to \mathbb{R}$ *be continuous on* $[a, b]$*, and be differentiable on* (a, b)*, and let* $g'(x) \neq 0$ *on* (a, b)*. If* $f'(x)/g'(x)$ *is increasing(deceasing) on* (a, b)*, then so are*

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ *is strictly monotone, then the monotonicity in the conclusion is also strict.*

From now on we let $r' = \sqrt{1 - r^2}$ for $0 < r < 1$.

Lemma 2.9. *Let* $c \in (0, 1]$ *and* $r \in (0, 1)$*.*

(1) The function $f_c(r) \equiv \frac{1-(cr')^2}{r\mathrm{arth}(cr)}$ *is strictly decreasing and concave with range* $(0, 1)$ *if* $c = 1$ *and strictly decreasing with range* $(0, \infty)$ *if* $0 < c < 1$*.*

(2) The function $F_c(r) \equiv \mathrm{arth}(cr)\mathrm{arth}(cr')$ *is strictly increasing on* $(0, \frac{\sqrt{2}}{2}]$ *and strictly decreasing on* $[\frac{\sqrt{2}}{2}, 1)$ *with maximum value* $(\mathrm{arth}(\frac{\sqrt{2}}{2}c))^2$*.*

Proof. (1) If $c = 1$, then $f_1(r) = \frac{r}{\mathrm{arth}\, r}$. By differentiation,

$$f_1'(r) = \frac{h_{11}(r)}{h_{12}(r)},$$

where $h_{11}(r) = \mathrm{arth} r - \frac{r}{r'^2}$ and $h_{12}(r) = (\mathrm{arth} r)^2$. It is easy to see that $h_{11}(0^+) = h_{12}(0^+) = 0$. Then

$$\frac{h_{11}'(r)}{h_{12}'(r)} = -\frac{h_{13}(r)}{h_{14}(r)},$$

where $h_{13}(r) = (\frac{r}{r'})^2$ and $h_{14}(r) = \mathrm{arth}\, r$. Then $h_{13}(0^+) = h_{14}(0^+) = 0$. By differentiation, we have

$$\frac{h_{13}'(r)}{h_{14}'(r)} = \frac{2r}{r'^2}$$

which is strictly increasing. Hence by Lemma 2.8, $\frac{h_{11}'(r)}{h_{12}'(r)}$ is strictly decreasing and so is f_1' with $f_1'(r) < f_1'(0^+) = 0$. Therefore, f_1 is strictly decreasing and concave on $(0, 1)$. The limiting value $f_1(1^-) = 0$ is clear and $f_1(0^+) = 1$ by l'Hôpital's Rule.

If $0 < c < 1$, then

$$f_c(r) = f_1(cr)h(r),$$

where $h(r) = \frac{1-(cr')^2}{cr^2}$. By differentiation,

$$h'(r) = -\frac{2(1 - c^2)}{cr^3} < 0.$$

Therefore, f_c is strictly decreasing. The limiting values are clear.

(2) By differentiation,

$$F_c'(r)=\frac{c}{r'f_c(r)}\left(\frac{f_c(r)}{f_c(r')}-1\right).$$

$\frac{f_c(r)}{f_c(r')}$ is strictly decreasing from $(0,1)$ onto $(0,\infty)$ by (1) and $F_c'(\frac{\sqrt{2}}{2})=0$. Then F_c is strictly increasing on $(0,\frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2},1)$ with maximum value $F_c(\frac{\sqrt{2}}{2})$. $\square$

Lemma 2.10. *Let* $c\in(0,1]$, $r\in(0,1)$, $m=\sqrt{(2-c^2)(3c^2-2)}$ *and* $r_0=\sqrt{\frac{1-m/c^2}{2}}$. *Let*

$$G_c(r)\equiv \operatorname{arth}(cr)+\operatorname{arth}(cr').$$

(1) If $0<c\le\sqrt{\frac{2}{3}}$, *then the range of* G_c *is* $(\operatorname{arth}c,\operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2})]$.

(2) If $\sqrt{\frac{2}{3}}<c<\sqrt{2(\sqrt{2}-1)}$, *then the range of* G_c *is* $(\operatorname{arth}c,\operatorname{arth}(\frac{c(r_0+r_0')}{1+c^2r_0r_0'})]$.

(3) If $\sqrt{2(\sqrt{2}-1)}\le c<1$, *then the range of* G_c *is* $[\operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2}),\operatorname{arth}(\frac{c(r_0+r_0')}{1+c^2r_0r_0'})]$.

(4) If $c=1$, *then the range of* G_c *is* $[\operatorname{arth}(\frac{2\sqrt{2}}{3}),\infty)$.

Proof. It is clear that the limiting values

$$G_c(0^+)=G_c(1^-)=\begin{cases}\operatorname{arth}c, & 0<c<1,\\ \infty, & c=1.\end{cases}$$

Let

$$g_c(r)=\operatorname{th}G_c(r)=\frac{c(r+r')}{1+c^2rr'}.$$

By differentiation, we have

$$r'(1+c^2rr')^2g_c'(r)=c(r'-r)(1-c^2-c^2rr').$$

Making substitution of $x=r^2$, we get

$$\begin{aligned}&1-c^2-c^2rr'=0\\ \Leftrightarrow\quad &c^4x^2-c^4x+(1-c^2)^2=0.\end{aligned}\tag{2.11}$$

Therefore, equation (2.11) has no root if $0<c^2<\frac{2}{3}$ or $c=1$, only one root $\frac{\sqrt{2}}{2}$ if $c^2=\frac{2}{3}$ and two different roots $r_0\,,r_0'$ if $\frac{2}{3}<c^2<1$. It is obvious that $r_0\in(0,\frac{\sqrt{2}}{2})$.

It is easy to see that $c=g_c(0)<g_c(\frac{\sqrt{2}}{2})=\frac{2\sqrt{2}}{2+c^2}c$ if and only if $c^2<2(\sqrt{2}-1)\approx0.828427$. If $\sqrt{\frac{2}{3}}<c<1$, g_c is increasing on $(0,r_0)$, $(\frac{\sqrt{2}}{2},r_0')$ and decreasing on $(r_0,\frac{\sqrt{2}}{2})$, $(r_0',1)$.

Now we get the following results.

(1) If $0<c\le\sqrt{\frac{2}{3}}$, $g_c'(r)=0$ is equivalent to $r=\frac{\sqrt{2}}{2}$. Then

$$\operatorname{arth}c=\operatorname{arth}(g_c(0))<G_c(r)\le\operatorname{arth}(g_c(\frac{\sqrt{2}}{2}))=\operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2}).$$

(2) If $\sqrt{\frac{2}{3}}<c<\sqrt{2(\sqrt{2}-1)}$, then

$$\operatorname{arth}c=\operatorname{arth}(g_c(0))<G_c(r)\le\operatorname{arth}(g_c(r_0))=\operatorname{arth}(\frac{c(r_0+r_0')}{1+c^2r_0r_0'}).$$

(3) If $\sqrt{2(\sqrt{2}-1)} \le c < 1$, then

$$\operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2}) = \operatorname{arth}(g_c(\frac{\sqrt{2}}{2})) \le G_c(r) \le \operatorname{arth}(g_c(r_0)) = \operatorname{arth}(\frac{c(r_0+r_0')}{1+c^2 r_0 r_0'}).$$

(4) If $c=1$, then

$$G_c(r) \ge \operatorname{arth}(g_1(\frac{\sqrt{2}}{2})) = \operatorname{arth}(\frac{2\sqrt{2}}{3}).$$

This completes the proof. □

Lemma 2.12. *Let $r \in (0,1)$.*

(1) The function $h_1(r) \equiv \frac{r'}{\operatorname{arth} r'}$ is strictly increasing and concave with range $(0,1)$.

(2) The function $h(r) \equiv \frac{r}{\operatorname{arth} r} + \frac{r'}{\operatorname{arth} r'}$ is strictly increasing on $(0, \frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2}, 1)$ and concave on $(0,1)$ with range $(1, \frac{\sqrt{2}}{\log(\sqrt{2}+1)}]$.

Proof. (1) The monotonicity and the limiting values of h_1 can be easily obtained by Lemma 2.9(1).

Now we prove the concavity of h_1. By differentiation,

$$h_1'(r) = \frac{r' - r^2 \operatorname{arth} r'}{r r' (\operatorname{arth} r')^2}$$

and

$$h_1''(r) r^2 r'^3 (\operatorname{arth} r')^3 = 2r'^2 - r' \operatorname{arth} r' - r^2 (\operatorname{arth} r')^2 \equiv \psi_1(r'). \tag{2.13}$$

Then $\psi_1(r) = 2r^2 - r \operatorname{arth} r - r'^2 (\operatorname{arth} r)^2$ and by differentiation

$$\psi_1'(r) = r(4 - \psi_2(r)),$$

where $\psi_2(r) = \frac{1}{r'^2} + 3\frac{\operatorname{arth} r}{r} - 2(\operatorname{arth} r)^2$. Since

$$\operatorname{arth} r = \frac{1}{2}\log\frac{1+r}{1-r} = \sum_{n=0}^{\infty} \frac{r^{2n+1}}{2n+1},$$

we have

$$\begin{aligned} r^2 r'^4 \psi_2'(r) &= (r^4 + 2r^2 - 3)\operatorname{arth} r - r^3 + 3r \\ &= \sum_{n=0}^{\infty} \frac{r^{2n+5}}{2n+1} + 2\sum_{n=1}^{\infty} \frac{r^{2n+3}}{2n+1} - 3\sum_{n=2}^{\infty} \frac{r^{2n+1}}{2n+1} \\ &= \sum_{n=2}^{\infty} \frac{16(n-1)}{(2n-3)(2n-1)(2n+1)} r^{2n+1} \\ &> 0. \end{aligned}$$

Hence ψ_2 is increasing with $\psi_2(0^+) = 4$ and ψ_1 is decreasing with $\psi_1(0^+) = 0$. Therefore by (2.13) h_1'' is negative and h_1' is decreasing. Then h_1 is concave on $(0,1)$.

(2) By differentiation,

$$h'(r) = f_1'(r) + h_1'(r) = f_1'(r) - \frac{r}{r'} f_1'(r'),$$

where $f_1(r)=\frac{r}{\operatorname{arth} r}=h_1(r')$. It is easy to see that $h'(\frac{\sqrt{2}}{2})=0$. Therefore, h is strictly increasing on $(0,\frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2},1)$. By (1) and Lemma 2.9(1), h' is strictly decreasing and $h(0^+)=h(1^-)=1$. Hence h is concave and

$$1<h(r)\le h(\frac{\sqrt{2}}{2})=\frac{\sqrt{2}}{\log(\sqrt{2}+1)}.$$

This completes the proof. $\square$

Lemma 2.14. *Let* $C=1-\frac{\log(\sqrt{2}+1)}{\sqrt{2}}\approx 0.376775$ *and* $r\in(0,1)$. *Let*

$$g(r)=\frac{r}{r'}(\frac{\operatorname{arth} r}{\operatorname{arth} r'})^{p-1}.$$

(1) g *is strictly decreasing if* $p\le 0$ *and strictly increasing if* $p\ge C$.

(2) If $p\in(0,C)$, *then there exists exactly one point* $r_0\in(0,\frac{\sqrt{2}}{2})$ *such that* g *is increasing on* $(0,r_0)$, $(r_0',1)$ *and decreasing on* (r_0,r_0').

(3) If $p\in(0,C)$, *then* $g(0^+)=0$ *and* $g(1^-)=\infty$.

Proof. (1) By logarithmic differentiation,

$$\begin{aligned}\frac{rr'^2\operatorname{arth} r\operatorname{arth} r'}{r\operatorname{arth} r'+r'\operatorname{arth} r}\cdot\frac{g'(r)}{g(r)} &= p-1+\frac{\operatorname{arth} r\operatorname{arth} r'}{r\operatorname{arth} r'+r'\operatorname{arth} r}\\ &= p-[1-\frac{1}{h(r)}],\end{aligned}$$

where $h(r)$ is as in Lemma 2.12(2). Since

$$0<1-\frac{1}{h(r)}\le C,$$

we see that g is strictly increasing if $p\ge C$ and decreasing if $p\le 0$.

(2) If $p\in(0,C)$, then there exists exactly one point $r_0\in(0,\frac{\sqrt{2}}{2})$ such that $g'(r_0)=g'(r_0')=0$ because $1-\frac{1}{h(r)}$ is increasing on $(0,\frac{\sqrt{2}}{2})$ and decreasing on $(\frac{\sqrt{2}}{2},1)$ by Lemma 2.12(2). Therefore, $g'>0$ if $r\in(0,r_0)\cup(r_0',1)$ and $g'<0$ if $r\in(r_0,r_0')$. Hence g is increasing on $(0,r_0)$, $(r_0',1)$ and decreasing on (r_0,r_0').

(3) Since $0<p<C<1$ and

$$\lim_{r\to 0^+}\frac{(\operatorname{arth} r')^{1-p}}{r^{-p}}=\lim_{r\to 0^+}\frac{1-p}{p}\cdot\frac{r^p}{r'(\operatorname{arth} r')^p}=0,$$

we have

$$\lim_{r\to 0^+}g(r)=\lim_{r\to 0^+}(\frac{r}{\operatorname{arth} r})^{1-p}\cdot(\frac{1}{r'})\cdot\frac{(\operatorname{arth} r')^{1-p}}{r^{-p}}=0$$

and

$$\lim_{r\to 1^-}g(r)=\lim_{r\to 1^-}(\frac{\operatorname{arth} r'}{r'})^{1-p}\cdot r\cdot\frac{r'^{-p}}{(\operatorname{arth} r)^{1-p}}=\infty.$$

This completes the proof. $\square$

Theorem 2.15. *Let* C *be as in Lemma 2.14. Then for all* $r\in(0,1)$,

$$H_p(\operatorname{arth} r,\operatorname{arth} r')\le\operatorname{arth}(\frac{\sqrt{2}}{2})\tag{2.16}$$

holds if and only if $p \le 0$ *and*

$$H_p(\operatorname{arth} r, \operatorname{arth} r') \ge \operatorname{arth}(\frac{\sqrt{2}}{2}) \tag{2.17}$$

holds if and only if $p \ge C$. *Equalities hold if and only if* $r = r' = \frac{\sqrt{2}}{2}$ *and all inequalities are sharp in both cases.*

Proof. We obtain the inequality (2.16) immediately if $p = 0$ by Lemma 2.9(2). Therefore, it suffices to discuss the case $p \neq 0$.

Let

$$f(r) = \frac{1}{p} \log \frac{(\operatorname{arth} r)^p + (\operatorname{arth} r')^p}{2}, \ p \neq 0.$$

By differentiation,

$$f'(r) = \frac{(\operatorname{arth} r')^{p-1}}{rr'[(\operatorname{arth} r)^p + (\operatorname{arth} r')^p]}[g(r) - 1],$$

where $g(r)$ is as in Lemma 2.14 and it is easy to see that $g(\frac{\sqrt{2}}{2}) = 1$.

Case 1. $p < 0$.

f is strictly increasing on $(0, \frac{\sqrt{2}}{2})$ and strictly decreasing on $(\frac{\sqrt{2}}{2}, 1)$ by Lemma 2.14(1). Therefore, $f(r) \le f(\frac{\sqrt{2}}{2})$ for all $p < 0$.

Case 2. $p \ge C$.

By Lemma 2.14(1) f is strictly decreasing on $(0, \frac{\sqrt{2}}{2})$ and strictly increasing on $(\frac{\sqrt{2}}{2}, 1)$ and hence $f(r) \ge f(\frac{\sqrt{2}}{2})$.

Case 3. $p \in (0, C)$.

By Lemma 2.14(2), there exists exactly one point $r_1 \in (0, r_0)$ such that $g(r_1) = g(r_1') = 1$, where $r_0 \in (0, \frac{\sqrt{2}}{2})$ is as in Lemma 2.14(2). Then f is strictly decreasing on $(0, r_1)$, $(\frac{\sqrt{2}}{2}, r_1')$ and strictly increasing on $(r_1, \frac{\sqrt{2}}{2})$, $(r_1', 1)$. Thus,

$$f(r_1) = f(r_1') < f(\frac{\sqrt{2}}{2}).$$

Since $f(0^+) = f(1^-) = \infty$, there exists $r_2 \in (0, r_1) \cup (r_1', 1)$ such that

$$f(\frac{\sqrt{2}}{2}) < f(r_2).$$

Therefore, inequalities (2.16) or (2.17) can not hold for all $r \in (0, 1)$.

This completes the proof of Theorem 2.15. ☐

Lemma 2.18. *Let* $r \in (0, 1)$.

(1) The function $f(r) \equiv \frac{r'^4 \operatorname{arth} r - r(1+r^2)}{r'^2[(1+r^2)\operatorname{arth} r - r]}$ *is strictly decreasing with range* $(-\infty, -2)$.

(2) For $p \in \mathbb{R}$ *define*

$$h_p(r) \equiv 1 + pr'^2 \frac{\operatorname{arth} r}{r} - (1 + r^2)\frac{\operatorname{arth} r}{r}.$$

(i) If $p \ge -2$, *then the range of* h_p *is* $(-\infty, p)$.

(ii) If $p < -2$, *then the range of* h_p *is* $(-\infty, C(p)]$, *where* $C(p) \equiv \sup_{0<r<1} h_p(r) \in (p\,, -1)$ *and* $\lim_{p \to -2} C(p) = -2$.

Proof. (1) Let $f_1(r) = r'^2\mathrm{arth}r - \frac{r(1+r^2)}{r'^2}$ and $f_2(r) = (1+r^2)\mathrm{arth}r - r$, then $f_1(0^+) = f_2(0^+) = 0$. By differentiation, we have

$$\frac{f_1'(r)}{f_2'(r)} = -1 - \frac{2r}{r'^2(r + r'^2\mathrm{arth}\,r)} = -1 - \frac{2}{r'^2(1+\frac{f_3(r)}{f_4(r)})},$$

where $f_3(r) = \mathrm{arth}\,r$ and $f_4(r) = \frac{r}{r'^2}$. It is easy to see that $f_3(0^+) = f_4(0^+) = 0$, then

$$\frac{f_3'(r)}{f_4'(r)} = \frac{r'^2}{1+r^2}$$

which is strictly decreasing. Hence by Lemma 2.8, $\frac{f_1'(r)}{f_2'(r)}$ is strictly decreasing and so is f with $f(0^+) = -2$. Since $(1+r^2)\mathrm{arth}r - r > 0$ and by l'Hôpital's Rule,

$$\lim_{r\to1^-}\frac{\mathrm{arth}r}{r'^{-2}} = \lim_{r\to1^-}\frac{r'^2}{2r} = 0, \tag{2.19}$$

we get $f(1^-) = -\infty$.

(2) By Lemma 2.12(1) and (2.19), it is easy to see that $h_p(0^+) = p$ and $h_p(1^-) = -\infty$. Next by differentiation, we have

$$h_p'(r) = \frac{1}{r}[(1+r^2)\frac{\mathrm{arth}r}{r} - 1][f(r) - p],$$

where $f(r)$ is as in (1).

If $p \geq -2$, then by (1) we see that $p > f(r)$ which implies that h_p is strictly decreasing and hence $h_p(r) < p$.

If $p < -2$, since the range of f is $(-\infty, -2)$, we see that there exists exactly one point $r_0 \in (0,1)$ such that $p = f(r_0)$. Then h_p is increasing on $(0, r_0)$ and decreasing on $(r_0, 1)$. Since

$$h_p(r) = -1 + 2(1 - \frac{\mathrm{arth}\,r}{r}) + (p+1)r'^2\frac{\mathrm{arth}\,r}{r} < -1,$$

by the continuity of h_p, there is a continuous function

$$C(p) \equiv \sup_{0<r<1} h_p(r)$$

with $p < C(p) < -1$ and $\lim_{p\to-2} C(p) = -2$. ☐

Lemma 2.20. *Let p, q be real numbers and $r \in (0,1)$. Let*

$$g_{p,q}(r) \equiv \frac{\mathrm{arth}^{q-1}r}{r^{p-1}r'^2}.$$

(1) If $p \geq -2$, then $g_{p,q}$ is strictly increasing for each $q \geq p$, and $g_{p,q}$ is not monotone for any $q < p$.
(2) If $p < -2$, then $g_{p,q}$ is strictly increasing for each $q \geq C(p)$, and $g_{p,q}$ is not monotone for any $q < C(p)$.

Here $C(p)$ is the same as in Lemma 2.18.

Proof. By logarithmic differentiation in r,

$$\frac{g_{p,q}'(r)}{g_{p,q}(r)} = \frac{1}{r'^2\mathrm{arth}r}(q - h_p(r)),$$

where $h_p(r)$ is the same as Lemma 2.18. Hence the results follow from Lemma 2.18(2). ☐

The following theorem studies the $H_{p,q}$-convexity of arth.

Theorem 2.21. *The inverse hyperbolic tangent function* arth *is strictly $H_{p,q}$-convex on* $(0,1)$ *if and only if* $(p,q) \in D_1 \cup D_2$*, where*

$$D_1 = \{(p,q)| -2 \leq p < +\infty, p \leq q < \infty\},$$

$$D_2 = \{(p,q)| -\infty < p \leq -2, C(p) \leq q < \infty\},$$

and $C(p)$ *is a continuous function with* $C(-2) = -2$*,* $C(-\infty) = -\infty$ *and* $p < C(p) < -1$ *for* $-\infty < p < -2$*. There are no values of* p *and* q *for which* arth *is* $H_{p,q}$*-concave on the whole interval* $(0,1)$*.*

Proof. The proof is divided into the following four cases.

Case 1. $p \neq 0$ and $q \neq 0$.

We may suppose that $0 < x \leq y < 1$. Define

$$F(x,y) = \mathrm{arth}^q\,(H_p(x,y)) - \frac{\mathrm{arth}^q x + \mathrm{arth}^q y}{2}.$$

Let $t = H_p(x,y)$, then $\frac{\partial t}{\partial x} = \frac{1}{2}(\frac{x}{t})^{p-1}$. If $x < y$, we see that that $t > x$. By differentiation, we have

$$\frac{\partial F}{\partial x} = \frac{q}{2} x^{p-1} \left(\frac{\mathrm{arth}^{q-1} t}{t^{p-1} t'^2} - \frac{\mathrm{arth}^{q-1} x}{x^{p-1} x'^2} \right).$$

Case 1.1. $p \geq -2$, $q \geq p$ and $pq \neq 0$. By Lemma 2.20(1), $\frac{\partial F}{\partial x} < 0$ if $p \leq q < 0$ and $\frac{\partial F}{\partial x} > 0$ if $q > 0$. Then $F(x,y)$ is strictly decreasing and $F(x,y) \geq F(y,y) = 0$ if $p \leq q < 0$, and $F(x,y)$ is strictly increasing and $F(x,y) \leq F(y,y) = 0$ if $q > 0$. Hence we have

$$\mathrm{arth}(H_p(x,y)) \leq H_q(\mathrm{arth} x, \mathrm{arth} y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$-convex on $(0,1)$ for $(p,q) \in \{(p,q)| -2 \leq p < 0, p \leq q < 0\} \cup \{(p,q)| -2 \leq p < 0, q > 0\} \cup \{(p,q)| 0 < p < +\infty, q \geq p\}$.

Case 1.2. $p \geq -2$, $q < p$ and $pq \neq 0$. By Lemma 2.20(1), with an argument similar to Case 1.1, it is easy to see that arth is neither $H_{p,q}$-concave nor $H_{p,q}$-convex on the whole interval $(0,1)$.

Case 1.3. $p < -2$, $q \geq C(p)$ and $pq \neq 0$. By Lemma 2.20(2), $\frac{\partial F}{\partial x} < 0$ if $C(p) \leq q < 0$ and $\frac{\partial F}{\partial x} > 0$ if $q > 0$. Then $F(x,y)$ is strictly decreasing and $F(x,y) \geq F(y,y) = 0$ if $p \leq q < 0$, and $F(x,y)$ is strictly increasing and $F(x,y) \leq F(y,y) = 0$ if $q > 0$. Hence we have

$$\mathrm{arth}(H_p(x,y)) \leq H_q(\mathrm{arth} x, \mathrm{arth} y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$-convex on $(0,1)$ for $(p,q) \in \{(p,q)|p < -2, C(p) \leq q < 0\} \cup \{(p,q)|p < -2, q > 0\}$.

Case 1.4. $p < -2$, $q < C(p)$. By Lemma 2.20(2), with an argument similar to Case 1.1, it is easy to see that arth is neither $H_{p,q}$-concave nor $H_{p,q}$-convex on the whole interval $(0,1)$.

Case 2. $p \neq 0$ and $q = 0$. For $0 < x \leq y < 1$, let

$$F(x,y) = \frac{\mathrm{arth}^2(H_p(x,y))}{\mathrm{arth} x \, \mathrm{arth} y},$$

and $t = H_p(x,y)$. If $x < y$, we see that that $t > x$. By logarithmic differentiation, we obtain

$$\frac{1}{F}\frac{\partial F}{\partial x} = x^{p-1} \left[\frac{(\mathrm{arth} t)^{-1}}{t^{p-1} t'^2} - \frac{(\mathrm{arth} x)^{-1}}{x^{p-1} x'^2} \right].$$

Case 2.1. $-2 \leq p < 0$ and $q = 0 > p$. By Lemma 2.20(1), we have $\frac{\partial F}{\partial x} > 0$ and $F(x,y) \leq F(y,y) = 1$. Hence we have

$$\mathrm{arth}(H_p(x,y)) \leq \sqrt{\mathrm{arth} x \, \mathrm{arth} y}$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$-convex on $(0,1)$ for $(p,q) \in \{(p,q) | -2 \leq p < 0, q = 0\}$.

Case 2.2. $p > 0$ and $q = 0 < p$. By Lemma 2.20(1), with an argument similar to Case 1.1, it is easy to see that arth is neither $H_{p,q}$-concave nor $H_{p,q}$-convex on the whole interval $(0,1)$.

Case 2.3. $p < -2$. We have $q = 0 \geq C(p)$, and by Lemma 2.20(2), with an argument similar to Case 1.1, it is easy to see that arth is $H_{p,q}$-convex on $(0,1)$ for $(p,q) \in \{(p,q) | p < -2, q = 0\}$.

Case 3. $p = 0$ and $q \neq 0$. For $0 < x \leq y < 1$, let

$$F(x,y) = \mathrm{arth}^q(\sqrt{xy}) - \frac{\mathrm{arth}^q x + \mathrm{arth}^q y}{2},$$

and $t = \sqrt{xy}$. If $x < y$, we have that $t > x$. By differentiation, we obtain

$$\frac{\partial F}{\partial x} = \frac{q}{2x} \left[\frac{\mathrm{arth}^{q-1} t}{t^{-1} t'^2} - \frac{\mathrm{arth}^{q-1} x}{x^{-1} x'^2} \right].$$

Case 3.1. $q > p = 0$. By Lemma 2.20(1), we have $\frac{\partial F}{\partial x} > 0$ and $F(x,y) \leq F(y,y) = 0$. Hence we have

$$\mathrm{arth}(\sqrt{xy}) \leq H_q(\mathrm{arth} x, \ \mathrm{arth} y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$-convex on $(0,1)$ for $(p,q) \in \{(p,q) | p = 0, q > 0\}$.

Case 3.2. $q < p = 0$. By Lemma 2.20(1), with an argument similar to Case 1.1, it is easy to see that arth is neither $H_{p,q}$-concave nor $H_{p,q}$-convex on the whole interval $(0,1)$.

Case 4. $p = q = 0$. By Case 1.1, for all $x\,, y \in (0,1)$, we have

$$\mathrm{arth}(H_p(x,y)) \leq H_p(\mathrm{arth} x, \ \mathrm{arth} y), \quad \text{for} \quad p \geq -2 \quad \text{and} \quad p \neq 0.$$

By the continuity of H_p in p and arth in x, we have

$$\mathrm{arth}(H_0(x,y)) \leq H_0(\mathrm{arth} x, \mathrm{arth} y).$$

In conclusion, arth is strictly $H_{0,0}$-convex on $(0,1)$.

This completes the proof of Theorem 2.21. □

Setting $p = 1 = q$ in Theorem 2.21, we obtain the convexity of arth easily.

Corollary 2.22. *The inverse hyperbolic tangent function* arth *is strictly convex on* $(0,1)$.

By (2.5), Theorem 2.21 has a simple application to the hyperbolic metric.

Corollary 2.23. *Let* $z \in S^1(H_p(|x|,|y|))$. *Then for all* $x, y \in \mathbb{B}^2 \setminus \{0\}$ *and* $p \geq -2$

$$\rho(0,z) \leq H_p(\rho(0,x), \rho(0,y)),$$

with equality if and only if $|x| = |y|$.

3. Some Propositions for Hyperbolic Metric

Next we give some geometric propositions for the hyperbolic metric.

Proposition 3.1. *Let c, d be arbitrary two points on the unit circle such that $0\,, c\,, d$ are noncollinear. Let $b \in [c,d] \cap \mathbb{B}^2$, $\{a\} = [0,b] \cap J^*[c,d]$, s be the midpoint of the Euclidean segment $[c,d]$. Then*
(1) a is the hyperbolic midpoint of the hyperbolic segment $J[0,b]$.
(2) s is on the hyperbolic circle $S_\rho(a,\rho(0,a)) = \{z|\rho(z,a) = \rho(0,a)\}$.



Figure 5. The point a is the hyperbolic midpoint of $J[0,b]$. The circular Arc (cad) is orthogonal to the unit circle. Here s is the Euclidean midpoint of the chord $[c,d]$.

Proof. Without loss of generality, we may assume $c = e^{i\alpha}$ and $d = e^{-i\alpha}$, where $0 < \alpha < \pi/2$.

(1) Since a is the hyperbolic midpoint of the hyperbolic segment $J[0,b]$, we have

$$\begin{aligned} & \rho(0,b) = 2\rho(0,a) \\ \Leftrightarrow\quad & \log\frac{1+|b|}{1-|b|} = 2\log\frac{1+|a|}{1-|a|} \\ \Leftrightarrow\quad & |b| = \frac{2|a|}{1+|a|^2}. \end{aligned} \tag{3.2}$$

Let $b = |b|e^{i\beta}(\,-\frac{\pi}{2} < \beta < \frac{\pi}{2})$. Then by the orthogonality of $J^*[c,d]$ and the unit circle

$$s = \cos\alpha = |b|\cos\beta = |b|\frac{|a|^2 + |w|^2 - r^2}{2|a||w|},$$

where $w = 1/\cos\alpha$ is the center and $r = \sqrt{|w|^2 - 1}$ is the radius of the corresponding orthogonal circle. Therefore, (3.2) holds and a is the hyperbolic midpoint of $J[0,b]$.

(2) It is easy to see that $[0,b]$ is also the Euclidean diameter of $S_\rho(a,\rho(0,a))$ by geometric observation and (1). Therefore, s is on the hyperbolic circle $S_\rho(a,\rho(0,a))$, see Figure 5. ☐

Proposition 3.3. *Let* $J_1 = J^*[e^{i\alpha}, -e^{-i\alpha}]$, $J_2 = J^*[-e^{i\alpha}, e^{-i\alpha}]$ *be two hyperbolic geodesics in* $\mathbb{B}^2$, $0 < \alpha < \pi/2$. *Let* $\{te_2\} = J_1 \cap Y$, *see Figure 6. Then*

$$d_\rho(J_1, J_2) = \rho(-te_2, te_2) = 2\log\frac{1+t}{1-t}. \tag{3.4}$$

Proof. Let $g : \mathbb{B}^2 \to \mathbb{H}^2$ be a Möbius transformation which satisfies $g(i) = \infty$ and $g(-i) = 0$. Then $g([-te_2, te_2]) = [g(-te_2), g(te_2)]$ which is on the ray emanating from 0 and perpendicular to ∂H^2. By the orthogonality of $J_i (i = 1, 2)$ and Y, we get

$$J_1^* = g(J_1) = \mathrm{Arc}(g(-e^{-i\alpha})g(te_2)g(e^{i\alpha}))\ \ ,\ \ J_2^* = g(J_2) = \mathrm{Arc}(g(-e^{i\alpha})g(-te_2)g(e^{-i\alpha})).$$

Here $g(-e^{-i\alpha}) = -g(e^{i\alpha})$ and $g(-e^{i\alpha}) = -g(e^{-i\alpha})$, see Figure 7. For every rectifiable arc γ_{xy}, $x \in J_1^*$ and $y \in J_2^*$, by geometric observation we get

$$\int_{\gamma_{xy}} \frac{1}{d(z, \partial\mathbb{H}^2)}|dz| \geq \int_{[g(-te_2), g(te_2)]} \frac{1}{d(z, \partial\mathbb{H}^2)}|dz|,$$

and hence

$$d_\rho(J_1^*, J_2^*) = \rho(g(-te_2), g(te_2)). \tag{3.5}$$

Since the hyperbolic distance is invariant under Möbius transformations, by (3.5) we get

$$d_\rho(J_1, J_2) = \rho(-te_2, te_2) = 2\log\frac{1+t}{1-t}.$$

□

Let $J_3 = J^*[e^{i\alpha}, e^{-i\alpha}]$ and $J_4 = J^*[-e^{-i\alpha}, -e^{i\alpha}]$. Let $\{se_1\} = J_3 \cap X$. Since the hyperbolic distance is invariant under rotations, by Proposition 3.3 we also obtain

$$d_\rho(J_3, J_4) = \rho(-se_1, se_1). \tag{3.6}$$

By the proof of Proposition 3.3, $J_3^* = g(J_3) = \mathrm{SArc}(g(e^{i\alpha})g(e^{-i\alpha}))$ and $J_4^* = g(J_4) = \mathrm{SArc}(g(-e^{-i\alpha})g(-e^{i\alpha}))$. Then g maps $\mathrm{Arc}(-i(se_1)i)$ to the ray emanating from 0 and tangent to J_3^* at point $g(se_1)$. Similarly, g maps $\mathrm{Arc}(-i(-se_1)i)$ to the ray emanating from 0 and tangent to J_4^* at point $g(-se_1)$. Then we have

$$d_\rho(J_3^*, J_4^*) = \rho(g(-se_1), g(se_1)). \tag{3.7}$$



FIGURE 6


H^2
J_1^*
$g(\mathrm{te}_2)$
d_2^*
d_1^*
J_2^*
$g(\mathrm{se}_1)$
J_4^*
J_3^*
$g(-e^{-\mathrm{i}\alpha})$
$g(-e^{\mathrm{i}\alpha})$
0
$g(e^{-\mathrm{i}\alpha})$
$g(e^{\mathrm{i}\alpha})$

FIGURE 7

The Möbius transformation g maps the unit disk in Figure 6 onto the upper half plane in Figure 7.

Remark 3.8. *By (3.4) and (3.6), we find the two distances between the opposite sides for the hyperbolic quadrilaterals with four vertices* $e^{i\alpha}$, $-e^{-i\alpha}$, $-e^{i\alpha}$, $e^{-i\alpha}$ *on the unit circle and the counterpart for the upper half plane by (3.5) and (3.7).*

4. Proof of Main Results

Proof of Theorem 1.1

Proof. Since the hyperbolic distance is Möbius invariant, we may assume that $v_a = 0$, v_b is on X, v_d is on Y and $v_c = te^{i\theta}$, $0 < t \le 1$ and $0 < \theta < \frac{\pi}{2}$(see Figure 1). Then by (2.7) the circle $S^1(b, r_b)$ through v_c and $\overline{v_c}$ is orthogonal to $\partial\mathbb{B}^2$, where

$$b = \frac{1+t^2}{2t\cos\theta} \quad \text{and} \quad r_b = \frac{\sqrt{(1+t^2)^2 - 4t^2\cos^2\theta}}{2t\cos\theta}.$$

By Proposition 3.3, we get

$$d_1 = \rho(0, v_b) = \log\frac{1+(b-r_b)}{1-(b-r_b)} = \text{arth}(\frac{2t}{1+t^2}\cos\theta).$$

Similarly, we get

$$d_2 = \rho(0, v_d) = \text{arth}(\frac{2t}{1+t^2}\sin\theta).$$

Then

$$d_1 d_2 = \text{arth}(Cr)\text{arth}(Cr'),$$

where $C = \frac{2t}{1+t^2} = \text{th}\rho(0, v_c) \in (0, 1]$ and $r = \cos\theta \in (0, 1)$. By Lemma 2.9(2), we have

$$d_1 d_2 \le (\text{arth}(\frac{\sqrt{2}}{2}C))^2.$$

This completes the proof of Theorem 1.1. ☐

Proof of Theorem 1.2

Proof. By the proof of Theorem 1.1, we have

$$d_1 + d_2 = \text{arth}(Cr) + \text{arth}(Cr'),$$

where $C \in (0, 1]$ and $r \in (0, 1)$. Then by Lemma 2.10, the desired conclusion follows. ☐

Proof of Corollary 1.4

Proof. There exists a Möbius transformation g which maps a, b, c, d to $e^{i\alpha}$, $-e^{-i\alpha}$, $-e^{i\alpha}$, $e^{-i\alpha}$, respectively, where $\alpha = \arccos\sqrt{1/|a, b, c, d|} \in (0\,, \pi/2)$, see Figure 6. By Proposition 3.3 and the proof of Theorem 1.1, we have

$$d_1 = 2\text{arth}r \quad \text{and} \quad d_2 = 2\text{arth}r' \tag{4.1}$$

where $r = \cos\alpha$.

Therefore, by Theorem 1.1 and Theorem 1.2, we have

$$d_1 d_2 = 4(\text{arth}\, r)(\text{arth}\, r') \le (2\text{arth}\,(\frac{\sqrt{2}}{2}))^2$$

and

$$d_1 + d_2 = 2(\text{arth}\, r + \text{arth}\, r') \ge 4\text{arth}\,(\frac{\sqrt{2}}{2}).$$

Equalities hold if and only if $\alpha = \frac{\pi}{4}$, namely, $|a, b, c, d| = \frac{1}{\cos^2\alpha} = 2$.

This completes the proof of Corollary 1.4. ☐

For $r \in (0,1)$ and $K \geq 1$ we define the distortion function

$$\varphi_K(r) = \mu^{-1}(\mu(r)/K)\,,$$

where $\mu(r)$ is the modulus of the planar Grötzsch ring, see [V, p.68, Exercise 5.61].

Lemma 4.2. [BV, Theorem 1.10] *Let $f : \mathbb{B}^2 \to \mathbb{B}^2$ be a K-quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$, and let ρ be the hyperbolic metric of $\mathbb{B}^2$. Then*

$$\rho(f(x), f(y)) \leq c(K) \max\{\rho(x,y), \rho(x,y)^{1/K}\}$$

for all $x, y \in \mathbb{B}^2$, where $c(K) = 2\mathrm{arth}(\varphi_K(\mathrm{th}\frac{1}{2}))$ and

$$K \leq u(K-1)+1 \leq \log(\mathrm{ch}(K\mathrm{arch}(e))) \leq c(K) \leq v(K-1)+K$$

with $u = \mathrm{arch}(e)\mathrm{th}(\mathrm{arch}(e)) > 1.5412$ and $v = \log(2(1+\sqrt{1-\frac{1}{e^2}})) < 1.3507$. In particular, $c(1) = 1$.

Proof of Theorem 1.6

Proof. In the same way as in the proof of Theorem 1.1, we still assume that $v_a = 0$, v_b is on X, v_d is on Y and $v_c = te^{i\theta}$, $0 < t \leq 1$ and $0 < \theta < \frac{\pi}{2}$(see Figure 1). Then

$$d_1 = \rho(0, v_b) = \mathrm{arth}(Cr) \;\text{ and }\; d_2 = \rho(0, v_d) = \mathrm{arth}(Cr'),$$

where $0 < C \leq 1$ and $0 < r < 1$.

By Lemma 4.2, we have

$$\begin{aligned} D_1D_2 &\leq \rho(f(0), f(v_b))\rho(f(0), f(v_d)) \\ &\leq c(K)^2 \max\{\rho(0,v_b), \rho(0,v_b)^{1/K}\} \cdot \max\{\rho(0,v_d), \rho(0,v_d)^{1/K}\} \\ &= c(K)^2 \max\{d_1, d_1^{1/K}\} \cdot \max\{d_2, d_2^{1/K}\}. \end{aligned}$$

(1) $0 < C \leq \frac{e^2-1}{e^2+1} \approx 0.761594$.

This implies that $d_1 < 1$ and $d_2 < 1$. Then by Theorem 1.1,

$$D_1D_2 \leq c(K)^2(d_1d_2)^{1/K} \leq c(K)^2(\mathrm{arth}(\frac{\sqrt{2}}{2}C))^{2/K}.$$

(2) $\frac{e^2-1}{e^2+1} < C \leq 1$.

Case 1. $\frac{0.761594}{C} \approx \frac{1}{C}\frac{e^2-1}{e^2+1} = r_C < r < 1$.

This implies that $d_1 > 1$ and $d_2 < 1$. Then

$$d_1 d_2^{1/K} = \mathrm{arth}(Cr)(\mathrm{arth}(Cr'))^{1/K} \equiv F_{C,K}(r).$$

By logarithmic differentiation, we have

$$\frac{F'_{C,K}(r)}{F_{C,K}(r)} = \frac{Cr}{r'(1-(Cr')^2)\mathrm{arth}(Cr')}\left[\frac{f_C(r)}{f_C(r')} - \frac{1}{K}\right], \tag{4.3}$$

where $f_C(r) = \frac{1-(Cr')^2}{r\mathrm{arth}(Cr)}$. By Lemma 2.9(1), $\frac{f_C(r)}{f_C(r')}$ is strictly decreasing from $(r_C, 1)$ onto $(0, \frac{1}{M_C})$. Here $M_C = \frac{f_C(r'_C)}{f_C(r_C)} > 1$ since $r_C \approx \frac{0.761594}{C} \geq 0.761594 > 0.707107 \approx \frac{\sqrt{2}}{2}$.

Case 1.1 $1 \leq K \leq M_C$. By (4.3), we have $F'_{C,K}(r) \leq 0$ and hence $F_{C,K}(r)$ is strictly decreasing on $(r_C, 1)$. Therefore,

$$d_1 d_2^{1/K} \leq F_{C,K}(r_C) = \mathrm{arth}(Cr_C)(\mathrm{arth}(Cr'_C))^{1/K}. \tag{4.4}$$

Case 1.2 $K > M_C$. There exists exactly one point $r_C(K) \in (r_C, 1)$ such that $\frac{f_C(r_C(K))}{f_C(\sqrt{1-r_C(K)^2})} = \frac{1}{K}$. Then $F_{K,C}$ is strictly increasing on $(r_C, r_C(K))$ and strictly decreasing on $(r_C(K), 1)$. Therefore,

$$d_1 d_2^{1/K} \le F_{K,C}(r_C(K)) = \mathrm{arth}(Cr_C(K))(\mathrm{arth}(C\sqrt{1-r_C(K)^2}))^{1/K}. \tag{4.5}$$

Case 2. $\sqrt{1-r_C^2} < r < r_C$.

This implies that $d_1 < 1$ and $d_2 < 1$. Since $\frac{\sqrt{2}}{2} \approx 0.707107 \in (\sqrt{1-r_C^2}, r_C)$, by Theorem 1.1 we have

$$(d_1 d_2)^{1/K} \le (\mathrm{arth}(\frac{\sqrt{2}}{2}C))^{2/K}. \tag{4.6}$$

Case 3. $0 < r < \sqrt{1-r_C^2}$.

This implies that $d_1 < 1$ and $d_2 > 1$. Putting $p = r'$, we have $r_C < p < 1$ and

$$d_1^{1/K} d_2 = \mathrm{arth}(Cp)(\mathrm{arth}(Cp'))^{1/K}.$$

Hence Case 3 is the same as Case 1.

Therefore, by (4.4) and (4.6), we have if $1 \le K \le M_C$,

$$D_1 D_2 \le c(K)^2 \max\{\mathrm{arth}(Cr_C)(\mathrm{arth}(C\sqrt{1-r_C^2}))^{1/K}, (\mathrm{arth}(\frac{\sqrt{2}}{2}C))^{2/K}\}.$$

And by (4.5) and (4.6), we have if $K > M_C$,

$$D_1 D_2 \le c(K)^2 \max\{\mathrm{arth}(Cr_C(K))(\mathrm{arth}(C\sqrt{1-r_C(K)^2}))^{1/K}, (\mathrm{arth}(\frac{\sqrt{2}}{2}C))^{2/K}\}.$$

This completes the proof of Theorem 1.6. ☐

Proof of Corollary 1.7

Proof. First, let g be the same as in the proof of Corollary 1.4. Let $\{se_1\} = g(J^*[a,d]) \cap X$, $\{te_2\} = g(J^*[a,b]) \cap Y$ and denote

$$z_1 = g^{-1}(se_1),\ z_2 = g^{-1}(te_2),\ z_3 = g^{-1}(-se_1),\ z_4 = g^{-1}(-te_2).$$

Since the hyperbolic distance is Möbius invariant, by the proof of Corollary 1.4, we get

$$d_1 = \rho(z_1, z_3) = 2\mathrm{arth}\, r \ \text{ and } \ d_2 = \rho(z_2, z_4) = 2\mathrm{arth}\, r',\ \ 0 < r < 1.$$

Then by Lemma 4.2, we have

$$\begin{aligned} D_1 D_2 &\le \rho(f(z_1), f(z_3))\rho(f(z_2), f(z_4)) \\ &\le c(K)^2 \max\{\rho(z_1, z_3), \rho(z_1, z_3)^{1/K}\} \cdot \max\{\rho(z_2, z_4), \rho(z_2, z_4)^{1/K}\} \\ &= c(K)^2 \max\{d_1, d_1^{1/K}\} \cdot \max\{d_2, d_2^{1/K}\}. \end{aligned}$$

Case 1. $0.886819 \approx \frac{2\sqrt{e}}{e+1} = r_1 < r < 1$.

This implies that $d_1 > 1$ and $d_2 < 1$. Then

$$d_1 d_2^{1/K} = 2^{1+1/K}\mathrm{arth}\, r(\mathrm{arth}\, r')^{1/K} \equiv 2^{1+1/K} F_{1,K}(r).$$

Let $M_1 = \frac{f_1(r_1')}{f_1(r_1)}$, where $f_1(r) = \frac{r}{\mathrm{arth} r}$. By the proof of Case 1 in Theorem 1.6, we have the following results.

Case 1.1 $1 \le K \le M_1$.

$$d_1 d_2^{1/K} \le 2^{1+1/K} F_{1,K}(r_1) = 2^{1+1/K}\mathrm{arth}\, r_1(\mathrm{arth}\, r_1')^{1/K}. \tag{4.7}$$

Case 2.2 $K > M_1$. There exists exactly one point $r_1(K) \in (r_1, 1)$ such that $\frac{f_1(r_1(K))}{f_1(\sqrt{1-r_1(K)^2})} = \frac{1}{K}$. Therefore,

$$d_1 d_2^{1/K} \le 2^{1+1/K} F_{1,K}(r_1(K)) = 2^{1+1/K} \mathrm{arth}\,(r_1(K))(\mathrm{arth}\,(\sqrt{1-r_1(K)^2}))^{1/K}. \tag{4.8}$$

Case 2. $0.462117 \approx \frac{e-1}{e+1} = \sqrt{1-r_1^2} < r < r_1 \approx 0.886819.$

This implies that $d_1 > 1$ and $d_2 > 1$. Since $\frac{\sqrt{2}}{2} \approx 0.707107 \in (\sqrt{1-r_1^2}, r_1)$, by Corollary 1.4 we have

$$d_1 d_2 \le \Big(2\log(\sqrt{2}+1)\Big)^2. \tag{4.9}$$

Case 3. $0 < r < \sqrt{1-r_1^2} \approx 0.462117.$

This implies that $d_1 < 1$ and $d_2 > 1$. Putting $p = r'$, we have $r_1 < p < 1$ and

$$d_1^{1/K} d_2 = 2^{1+1/K} \mathrm{arth}\, p(\mathrm{arth}\, p')^{1/K}.$$

Hence Case 3 is the same as Case 2.

Therefore, by (4.7) and (4.9), we have if $1 \le K \le M_1$,

$$D_1 D_2 \le c(K)^2 \max\{2^{1+1/K} \mathrm{arth}\, r_1 (\mathrm{arth}\, r_1')^{1/K}, \Big(2\log(\sqrt{2}+1)\Big)^2\}.$$

And by (4.8) and (4.9), we have if $K > M_1$,

$$D_1 D_2 \le c(K)^2 \max\{2^{1+1/K} \mathrm{arth}\,(r_1(K))(\mathrm{arth}\,(\sqrt{1-r_1(K)^2}))^{1/K}, \Big(2\log(\sqrt{2}+1)\Big)^2\}.$$

This completes the proof of Corollary 1.7. ☐

Acknowledgments. The research of Matti Vuorinen was supported by the Academy of Finland, Project 2600066611. The research of Gendi Wang was supported by CIMO of Finland, Grant TM-10-7364.

References

[AVV1] G.D. Anderson, M.K. Vamanamurthy, M. Vuorinen: *Conformal Invariants, Inequalities and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.

[AVV2] G.D. Anderson, M.K. Vamanamurthy, M. Vuorinen: *Generalized convexity and inequalities*, J. Math. Anal. Appl., 335(2007), 1294-1308.

[A] J.W. Anderson: *Hyperbolic Geometry*, 2nd Ed., Springer Undergraduate Mathematics Series, Springer-Verlag,London, 2005.

[Ba] Á. Baricz: *Convexity of the zero-balanced Gaussian hypergeometric functions with respect to Hölder means*, J. Inequal. Pure and Appl. Math., 8(2007)Art. 40, 9pp.

[Be] A. F. Beardon: *The Geometry of Discrete Groups*, Graduate Texts in Math. Vol. 91, Springer-Verlag, Berlin-Heidelberg-New York, 1982.

[BM] A. F. Beardon and D. Minda: *Conformal images and the angle of parallelism*, Comput. Methods Funct. Theory 12, 2012, 87–100.

[BV] B. A. Bhayo and M. Vuorinen: *On Mori's theorem for quasiconformal maps in the $n-$space*, Trans. Amer. Math. Soc. 363, 2011, 5703-5719.

[Bu] P.S. Bullen: *Handbook of Means and Their Inequalities*, Mathematics and Its Applications, vol. 560, Kluwer Academic Publishers, Dordrecht, 2003.

[CG] H. S. M. Coxeter and S. L. Greitzer: *Geometry Revisited*, The Math. Association of America, 1976.

[HLP] G. Hardy, J.E. Littlewood, G. Pólya: *Inequalities*, Cambridge Univ. Press, Cambridge, 1952.

[KL] L. Keen and N. Lakic: *Hyperbolic Geometry from a Local Viewpoint*, London Math. Soc. Student Texts 68, Cambridge Univ. Press, Cambridge, 2007.

[KV] R. Klén and M. Vuorinen: *Apollonian circles and hyperbolic geometry*, arXiv:1010.2630v2 [math. MG].

[K] J. G. Krzyz: Problems in complex variable theory. Translation of the 1962 Polish original. Modern Analytic and Computational Methods in Science and Mathematics, No.36. American Elsevier Publishing Co., Inc. New York; PWN-Polish Scientific Publishers, Warsaw, 1971, xvii+183 pp.

[M] J. Milnor: *Hyperbolic geometry: the first 150 years*, Bull. Amer. Math. Soc. (N. S.) 6, (1982), no.1, 9-24.

[V] M. Vuorinen: *Conformal Geometry and Quasiregular Mappings*, Lecture Notes in Math. 1319, Springer-Verlag, Berlin-Heidelberg, 1988.

[ZWC] X.H. Zhang, G.D. Wang, Y.M. Chu: *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl., 353(2009), 256-259.

Department of Mathematics and Statistics, University of Turku, Turku 20014, Finland
E-mail address: `vuorinen@utu.fi`
E-mail address: `genwan@utu.fi`